



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

PROBLEMS FOR SOLUTION.

ALGEBRA.

337. Proposed by I. M. CURTISS, Brooklyn, N. Y.

Three regiments move north as follows: B is 20 miles east of A; C is 20 miles south of B, and each marches 20 miles between the hours of 5 a. m. and 3 p. m. A horseman with a message from C starts at 5 a. m. and rides north till he overtakes B, then sets a straight course for the point at which he calculates to overtake A, then sets a straight course for the next point at which he will again overtake B, then rides south to the point where he first overtook B, reaching that point at the same time as C, namely 3 p. m. What uniform rate of travel enabled the messenger to do this?

Solution by G. B. M. ZERR, A. M., Ph. D., Philadelphia, Pa.

Let x = time required to overtake B. He travels $20 + 2x$ miles. Hence $\frac{20+2x}{x}$ = his rate. Let y = time to go from B to A. He travels $2\sqrt{100+y^2}$ miles. $\frac{2\sqrt{100+y^2}}{y}$ = his rate. After reaching B a second time he has left $10 - x - 2y$ hours to go $2x + 4y$ miles.

$\therefore \frac{2x+4y}{10-x-2y}$ = his rate. But his rate is uniform. Hence we get

$$\frac{20+2x}{x} = \frac{2\sqrt{100+y^2}}{y}, \text{ or } 5x^2 = 5y^2 + xy \dots (1).$$

$$\frac{20+2x}{x} = \frac{2x+4y}{10-x-2y}, \text{ or } x^2 + 2xy = 50 - 10y \dots (2).$$

If $x = vy$ in (1), we get $5v^2 - v = 5$ or $v = 1.10499$.

$\therefore x = 1.10499y$. This in (2) gives $14.41996y^2 + 10y = 50$ or $y = 1.54737$.
 $x = 1.70983$. Rate = $20/x + 2 = 13.69707$ miles an hour.

Also solved by V. M. Spunar, A. H. Holmes, and J. Scheffer.

338. Proposed by R. D. CARMICHAEL, Princeton University.

Prove that $\pi = 3 + \frac{1}{3} \cdot \frac{1}{1.2} - \frac{1}{5} \cdot \frac{1}{2.3} + \frac{1}{7} \cdot \frac{1}{3.4} - \frac{1}{9} \cdot \frac{1}{4.5} + \dots$

Solution by S. LEFSEHETZ, East Pittsburg, Pa.

$$\text{Let } S(x) = \sum_1^{\infty} (-1)^{n+1} \frac{x^{2n}}{n(n+1)(2n+1)} = \sum_1^{\infty} \left[\frac{1}{n} + \frac{1}{n+1} - \frac{4}{2n+1} \right] (-1)^{n+1} x^{2n}$$

$$= \sum_1^{\infty} (-1)^{n+1} \frac{x^{2n}}{n} + \sum_1^{\infty} (-1)^{n+1} \frac{x^{2n}}{n+1} + 4 \sum_1^{\infty} (-1)^2 \frac{x^{2n}}{2n+1}$$